



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

G , $FC=2BC=4r \sin \frac{\pi}{n}$ is the radius and $\frac{2\pi}{n}$ the central angle, from G to H ,
 $GD=3CD=6r \sin \frac{\pi}{n}$ is the radius and $\frac{2\pi}{n}$ the central angle and so on. For the
 n th side or the last side of the first revolution, radius $=2nr \sin \frac{\pi}{n}$, for the last
side of second revolution radius $=4nr \sin \frac{\pi}{n}$, for the last side of the
 m th revolution radius $=2mnr \sin \frac{\pi}{n}$.

Hence without the use of calculus if s = length of path, arc $AF=2r \times \frac{2\pi}{n} \sin \frac{\pi}{n}$, arc $FG=4r \times \frac{2\pi}{n} \sin \frac{\pi}{n}$, arc $GH=6r \times \frac{2\pi}{n} \sin \frac{\pi}{n}$.

$$\therefore s = 2r \times \frac{2\pi}{n} \sin \frac{\pi}{n} \{ 1 + 2 + 3 + \dots (mn-1) \} = 2r \times \frac{2\pi}{n} \times \sin \frac{\pi}{n} \frac{mn(mn-1)}{2}$$

$$= 2\pi rm(mn-1) \sin \frac{\pi}{n}. \quad \text{But } r=1, m=120.$$

$$\therefore s = 240\pi(120n-1) \sin \frac{\pi}{n}.$$

If $n=6$, $s=271057.248$ feet, if $n=\infty$ $s=2(120)^2 \pi^2 = 284245.9361$ feet.

Also solved by *ALFRED HUME* and the *PROPOSER*.

12. Proposed by *ISAAC L. BEVERAGE*, Monterey, Virginia.

Given the equations $2z^3 = x + 3z$ and $5z^2 = y + 2z$. To find $\frac{dy}{dx}$ for $x=0$.

Solution by Professor *P. H. PHILBRICK*, Lake Charles, Louisiana.

We have $x=2z^3-3z$, and $y=5z^2-2z$. $\therefore dx=(6z^2-3)dz$, $dy=(10z-2)dz$, and $\frac{dy}{dx} = \frac{10z-2}{6z^2-3}$.

But $x=2z^3-3z=z(2z^2-3)=0$. $\therefore z=0$, and $\pm \sqrt{\frac{3}{2}}$.

$\therefore \frac{dy}{dx} = \frac{2}{3}$, or $\pm \frac{5\sqrt{6}-2}{6}$, when $x=0$.

Also solved by Professors *BLACK*, *DRAUGHON*, *MATZ*, *SCHEFFER*, *WHITAKER*, and *ZERR*.

13. Proposed by *J. A. CALDERHEAD*, Superintendent of Schools, Lima, Ohio.

A steamer whose course is due west and speed 10 knots is sighted by another steamer going at 8 knots; what course must the latter steer, so as to cross the track of the former at the least possible distance from her?

Solution by *C. W. M. BLACK*, A. M., Department of Mathematics, Wilmington Conference Academy, Dover, Delaware.

Let A be the position of sighting steamer, D of other when sighted, B where A crosses course of D , which by that time has reached C . AE is $CE \perp$. Let $y=CB=CD+DB=CD+DE-BE$. Let a =no. hrs. A takes to reach B , and $b=AE$. $\angle EAB=x$, $\angle EAD=\theta$.

$$AE' = AB \cos x, \text{ or } b = 8a \cos x, a = \frac{b}{8 \cos x}. \quad CD = 10a.$$

$$DE = b \tan \theta, BE = b \tan x. \quad y = \frac{10b}{8 \cos x} + b \tan \theta - b \tan x. \quad \frac{dy}{dx} = \frac{5b \sin x}{4 \cos^2 x} - b \sec^2 x.$$

$$\text{Let } \frac{dy}{dx} = 0, \text{ whence } \sin \theta = \frac{4}{5}, \theta = 53^\circ 8'.$$

The equation shows that b and θ do not affect the result.

Also solved by *Professors PHILBRICK, WHITAKER, ZERR, and the PROPOSER.*

14. Proposed by J. F. W. SCHEFFER, A. M., Hagerstown, Maryland.

Right triangles are inscribed in a circle whose center $= (a, b)$, and radius $= c$. If one of the legs passes through a fixed point, prove that $c^2(x^2 + y^2) = (a^2 + b^2 - c^2 - ax - by)^2$ is the curve to which the other leg is always tangent; the fixed point being the origin of the co-ordinates.

Solution by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Let $lx + my = 1 \dots (1)$ be the chord of the circle whose envelope is to be found. Then $y = \frac{m}{l}x \dots (2)$ is the side of the triangle passing through the fixed point, and $(x-a)^2 + (y-b)^2 = c^2 \dots (3)$ is the equation to the circle.

$$(1) \text{ and } (2) \text{ intersect in } x' = \frac{l}{l^2 + m^2}, y' = \frac{m}{l^2 + m^2}.$$

This point being on (3), we must have on substituting x' and y' in (3), after reducing, $(a^2 + b^2 - c^2)(l^2 + m^2) - 2al - 2bm + 1 = 0 \dots (4)$. Making (4) homogeneous in l and m by means of (1) and arranging,

$$[x^2 - 2ax + (a^2 + b^2 - c^2)] \frac{l^2}{m^2} + 2(xy - bx - ay) \frac{l}{m} + [y^2 - 2by + (a^2 + b^2 - c^2)] = 0 \dots$$

..(5), a quadratic in the parameter $\frac{l}{m}$, giving the envelope

$$c^2(x^2 + y^2) = [(a^2 + b^2 - c^2) - by - ax]^2 \dots (6).$$

Also solved by P. S. BERG, G. B. M. ZERR and the PROPOSER.

15. Proposed by CHARLES E. MYERS, Canton, Ohio.

From a given quantity of material a cylindrical cup with circular bottom and open top is to be made, the cup to contain the greatest amount. What must be its dimensions?

Solution by F. P. MATZ, M. S., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

Represent the radius of the base by x , and the altitude by y ; then, obviously, $\pi x^2 + 2\pi xy = s$. $\therefore y = (s - \pi x^2) \div 2\pi x$.

$$\text{Also, } V = \pi x^2 y = \pi x^2 \left(\frac{s - \pi x^2}{2\pi x} \right), = a \text{ maximum.}$$

